
Exercise 6 for Theoretical Solid State Physics in Summer 2023

Prof. Dr. Andreas Klümper

Dr. Vladislav Popkov

Submission: 17.05.2023, 12:00 in the P.O. Box Popkov on D.10 (by e-mail)

Discussion: 17.05.2023, 14:15

1. Density of states (5 points)

In the lecture we have considered free particles with energy momentum dispersion $E(\vec{k}) = \vec{k}^2/2m$ in d -dimensions. We computed the density of states where we focused on $d = 1, 2, 3$.

Now let us assume that an energy momentum dispersion $E(\vec{k}) = v|\vec{k}|$ in d -dimensions is given. In other words, the energy is linear in the absolute value of the momentum \vec{k} .

- Compute the density of states and plot these for the cases $d = 1, 2, 3$.
- Which (fermionic) particles have a linear dispersion law?
- Have you heard about electrons on certain (2d) lattices with “linear regions” of the energy bands?

2. Van-Hove singularities (12 points)

The density of states $D_n(E)$ of the n -th energy band is given by

$$D_n(E) = \frac{1}{(2\pi)^3} \int_{E_n(\vec{k})=E} \frac{dS}{\left| \vec{\nabla}_{\vec{k}} E_n(\vec{k}) \right|}.$$

The total density of states is the sum over all band indices n . Analyse the different types of singularities. Expand the energies close to the critical point E_0

$$E_n(\vec{k}) = E_0 + \left(\frac{k_x^2}{2m_{xn}^*} + \frac{k_y^2}{2m_{yn}^*} + \frac{k_z^2}{2m_{zn}^*} \right) + \mathcal{O}(k^4)$$

and discuss the four different cases of relative signs of $m_{jn}^*, j = x, y, z$.